

Office of Naval Research  
Department of the Navy  
Contract Nonr 220(43)

A COMPUTER METHOD FOR CALCULATION  
OF THE COMPLETE AND INCOMPLETE  
ELLIPTIC INTEGRALS OF THE THIRD KIND

by

D. K. Ai and Z. L. Harrison

Hydrodynamics Laboratory  
Kármán Laboratory of Fluid Mechanics and Jet Propulsion  
California Institute of Technology  
Pasadena, California

Report No. E-110.3

February 1964

Office of Naval Research  
Department of the Navy  
Contract Nonr 220(43)

A COMPUTER METHOD FOR CALCULATION  
OF THE COMPLETE AND INCOMPLETE  
ELLIPTIC INTEGRALS OF THE THIRD KIND

by

D. K. Ai and Z. L. Harrison

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

Hydrodynamics Laboratory  
Karman Laboratory of Fluid Mechanics and Jet Propulsion  
California Institute of Technology  
Pasadena, California

Report No. E-110.3

Approved by: T. Y. Wu

A.J. Acosta

February 1964

## ABSTRACT

Numerical approximations and a Fortran IV program are given for the calculation by an IBM 7090 computer of the complete and incomplete elliptic integrals of the third kind. In its present form results are limited to six decimal places, but the method is valid for all values of amplitude  $\varphi$ , modulus  $k$  and real values of the parameter  $a^2$ . For the purpose of completeness, adaptations of other programs for complete and incomplete elliptic integrals of the first and second kind are also presented.

## TABLE OF CONTENTS

	Page
Abstract	
List of Symbols and Definitions	
1. Introduction	1
2. Complete Elliptic Integrals of the Third Kind	3
3. Incomplete Elliptic Integrals of the Third Kind	7
4. Appendices	
A. Approximate Functions for the Complete Elliptic Integrals of the First and Second Kind	17
B. Evaluation of the Incomplete Elliptic Integrals of the First and Second Kind	19
C. Flow Charts and Subroutines	24
References	27

## LIST OF SYMBOLS AND DEFINITIONS

$a^2$	parameter in the elliptic integral of the third kind.
$k, k'$	modulus and complementary modulus, $0 \leq k^2 \leq 1$ and $k' = \sqrt{1 - k^2}$ .
$y$ or $\varphi$	argument of the elliptic integral $0 < y \leq 1$ , $0 < \varphi \leq \frac{\pi}{2}$ .
$F(\varphi, k)$ , $E(\varphi, k)$ , $\Pi(\varphi, a^2, k)$	elliptic integrals of the first, second and third kind respectively, where

$$F(\varphi, k) = \int_0^y \frac{dt}{\sqrt{(1-t^2)(1-k^2 t^2)}} = \int_0^\varphi \frac{d\theta}{\sqrt{1-k^2 \sin^2 \theta}}$$

$$E(\varphi, k) = \int_0^y \sqrt{\frac{1-k^2 t^2}{1-t^2}} dt = \int_0^\varphi \sqrt{1-k^2 \sin^2 \theta} d\theta$$

and

$$\begin{aligned} \Pi(\varphi, a^2, k) &= \int_0^y \frac{1}{1-a^2 t^2} \frac{dt}{\sqrt{(1-t^2)(1-k^2 t^2)}} \\ &= \int_0^\varphi \frac{1}{1-a^2 \sin^2 \theta} \frac{d\theta}{\sqrt{1-k^2 \sin^2 \theta}} . \end{aligned}$$

$K, E$ and $\Pi(a^2, k)$	complete elliptic integrals of the first, second and third kind respectively, where
--------------------------	---

$$K = K(k) = \int_0^1 \frac{dt}{\sqrt{(1-t^2)(1-k^2 t^2)}} = \int_0^{\pi/2} \frac{d\theta}{\sqrt{1-k^2 \sin^2 \theta}}$$

$$E = E(k) = \int_0^1 \sqrt{\frac{1-k^2 t^2}{1-t^2}} dt = \int_0^{\pi/2} \sqrt{1-k^2 \sin^2 \theta} d\theta$$

$$\begin{aligned} \Pi(a^2, k) &= \int_0^1 \frac{1}{1-a^2 t^2} \frac{dt}{\sqrt{(1-t^2)(1-k^2 t^2)}} \\ &= \int_0^{\pi/2} \frac{1}{1-a^2 \sin^2 \theta} \frac{d\theta}{\sqrt{1-k^2 \sin^2 \theta}} . \end{aligned}$$

$K^*, E^*$  approximation functions of complete elliptic integrals of the first and second kind.

$Z(\varphi, k)$  Jacobian Zeta Function.

$$Z(\varphi, k) = E(\varphi, k) - \frac{E}{K} F(\varphi, k)$$

$\Lambda_o(\varphi, k)$  Heuman's Lambda Function.

$$\Lambda_o(\varphi, k) = \frac{2}{\pi} \{ E F(\varphi, k') + K [ E(\varphi, k') - F(\varphi, k') ] \}$$

## 1. Introduction

There are tables of the elliptic integral of the third kind, [1], [2]<sup>\*</sup>; these tables are somewhat tedious to use because of the limited number of entries and range necessary in a three parameter tabulation.<sup>\*\*</sup> Either cross plotting or non-linear interpolation are necessary to obtain values of  $\Pi$  within the range of the tables and furthermore, values of the integral outside the range of the tables must be calculated from cumbersome addition formulae as well as interpolation. These methods are undesirable when very many values of these integrals are needed and completely impractical for adaptation to computer use.

Our interest and need of evaluating these elliptic integrals arise in a study of free-streamline potential flows. This problem requires the numerical integration of expressions involving  $\Pi$  as well as evaluating all of the elliptic integrals many times. Since these calculations are performed on an IBM 7090 digital computer, it is necessary to have computer methods of evaluating these elliptic integrals as needed.

Three subroutines have been written in Fortran IV language to accomplish this result: the complete elliptic integrals of the first and second kind  $K$  and  $E$ ; the incomplete elliptic integrals of the first and second kind,  $F(\varphi, k)$  and  $E(\varphi, k)$ ; and most especially the complete and incomplete elliptic integrals of the third kind,  $\Pi(a^2, k)$  and  $\Pi(\varphi, a^2, k)$

---

\* The numbers in brackets refer to references at the end of the text.

\*\* In Ref. [1] for example,  $-1 < a^2 < 1$  and in Ref. [2]  $-1 < a^2 < 0$ .

respectively. The first two of these programs closely follow methods outlined by others, Refs. [3] and [4] and are detailed in Appendices A and B respectively.

The present programs are written in single precision arithmetic since that accuracy is sufficient for our calculations. The accuracy of the program for  $K$  and  $E$ , which uses the approximate formulae given in Ref. [3], would not be improved by converting to double precision arithmetic. The remaining subroutines use series representations and consequently their accuracy can be improved by carrying more significant figures. All programs are, in their present form, accurate to six figures with the following exceptions: when  $k \sin \varphi > 0.9$ ,  $E(\varphi, k)$  and  $F(\varphi, k)$  lose accuracy; when  $k^2 > 0.9$  or  $|a^2 - 1| < 0.1$ ,  $\Pi(\varphi, a^2, k)$  and  $\Pi(a^2, k)$  lose accuracy.\* Four figure accuracy is guaranteed except  $k \sin \varphi > 0.9999$  or  $|1 - a \sin \varphi| < 0.0001$ . Whenever the accuracy is less than six figures, comments are printed and indicators are set to inform the user of this loss in accuracy.

The flow chart and the Fortran IV program listings are included in Appendix C.

---

\* This result is because the accuracy in computing  $(1 - k \sin \varphi)$ ,  $(1 - a^2)$ , and  $(1 - a \sin \varphi)$  decreases as  $a$  and  $k$  approach unity as  $\varphi$  tends to  $\pi/2$ .



## 2. Complete Elliptic Integrals of the Third Kind

The complete elliptic integral of the third kind, i. e. ,

$$\Pi(a^2, k) = \int_0^1 \frac{1}{1-a^2 t^2} \frac{dt}{\sqrt{(1-t^2)(1-k^2 t^2)}} = \int_0^{\pi/2} \frac{1}{1-a^2 \sin^2 \theta} \frac{d\theta}{\sqrt{1-k^2 \sin^2 \theta}} \quad (1)$$

depends on the two parameters  $a^2$  and  $k$ . Considering only real values of  $a^2$ , the ranges of these parameters are

$$0 \leq k^2 \leq 1 \quad \text{and} \quad -\infty < a^2 < \infty .$$

Some special cases of  $a^2$  and  $k$  are as follows:

a)  $k = 0$ ; the integral can be integrated in closed form and is

$$\Pi(a^2, 0) = \begin{cases} \frac{\pi}{2} \frac{1}{\sqrt{1-a^2}} , & a^2 < 1 \\ \infty , & a^2 = 1 \\ 0^* , & a^2 > 1 \end{cases} \quad (2)$$

$$\text{b) } k = 1; \quad \Pi(a^2, 1) = \infty \quad (3)$$

$$\text{c) } a^2 = 0; \quad \Pi(0, k) = K \quad (4)$$

$$\text{d) } a^2 = \pm k; \quad \Pi(\pm k, k) = \frac{1}{4(1 \mp k)} [\pi + 2(1 \mp k) K] \quad (5)$$

$$\text{e) } a^2 = k^2; \quad \Pi(k^2, k) = \frac{E}{1-k^2} \quad (6)$$

$$\text{f) } a^2 = 1; \quad \Pi(1, k) = \infty \quad (7)$$

The remainder of the values of  $a^2$  and  $0 < k^2 < 1$  can be classified into four cases as shown in Fig. 1.

---

\* For  $a^2 > 1$ , the integral is interpreted by its Cauchy's principal value.

$$\begin{array}{ll}
 \text{Case I,} & 0 < -a^2 < \infty \\
 \text{Case II,} & k^2 < a^2 < 1
 \end{array}
 \left. \vphantom{\begin{array}{l} \text{Case I,} \\ \text{Case II,} \end{array}} \right\} \text{Circular cases;}$$

$$\begin{array}{ll}
 \text{Case III,} & 0 < a^2 < k^2 \\
 \text{Case IV,} & 1 < a^2 < \infty
 \end{array}
 \left. \vphantom{\begin{array}{l} \text{Case III,} \\ \text{Case IV,} \end{array}} \right\} \text{Hyperbolic cases;}$$




Figure 1

The integral,  $\Pi(a^2, k)$ , can be evaluated in terms of elementary functions and Heuman's Lambda function<sup>\*</sup>

$$\Lambda_0(\varphi, k) = \frac{2}{\pi} \{ E F(\varphi, k') + K [E(\varphi, k') - F(\varphi, k')] \} \quad (8)$$

for the circular cases (I and II) or of the Jacobian Zeta function

$$Z(\varphi, k) = E(\varphi, k) - \frac{E}{K} F(\varphi, k) \quad (9)$$

or  $K Z(\varphi, k) = K E(\varphi, k) - E F(\varphi, k) \quad (9a)$

for the Hyperbolic cases (III and IV).

<sup>\*</sup> These definitions and formulae can be found in Ref. [5].

The Lambda function and Zeta function as given in Eqs. (8) and (9a) are tabulated [5] or can readily be evaluated by using the methods described in Appendices A and B to calculate  $K$ ,  $E$ ,  $F(\varphi, k)$  and  $E(\varphi, k)$ .

Case I:  $0 < -a^2 < \infty$

$$\Pi(a^2, k) = \frac{k^2}{k^2 - a^2} K - \frac{\pi}{2} \frac{a^2 \Lambda_O(\psi, k)}{\sqrt{a^2(1-a^2)(a^2-k^2)}} \quad (10)$$

where

$$\psi = \sin^{-1} \sqrt{\frac{a^2}{a^2 - k^2}} \quad (11)$$

or

$$\Pi(a^2, k) = \frac{K}{1 - a^2} + \frac{\pi}{2} \frac{a^2 [\Lambda_O(\beta, k) - 1]}{\sqrt{a^2(1-a^2)(a^2-k^2)}} \quad (12)$$

where

$$\beta = \sin^{-1} \sqrt{\frac{1}{1 - a^2}} \quad (13)$$

It is obvious that less time is required to evaluate  $F(\varphi, k)$  and  $E(\varphi, k)$  when  $\varphi$  is small. Since

$$\psi < \beta \quad \text{when} \quad -k < a^2 < 0$$

and

$$\psi > \beta \quad \text{when} \quad -\infty < a^2 < -k$$

$\Pi(a^2, k)$  is programmed by Eq. (10) in the first interval and by Eq. (12) in the second interval.

Case II:  $k^2 < a^2 < 1$

$$\Pi(a^2, k) = K + \frac{\pi}{2} \frac{a[1 - \Lambda_O(\theta, k)]}{\sqrt{(a^2 - k^2)(1 - a^2)}} \quad (14)$$

where

$$\theta = \sin^{-1} \sqrt{\frac{1 - a^2}{1 - k^2}} \quad (15)$$

or

$$\Pi(a^2, k) = \frac{\pi}{2} \frac{a \Lambda_0(\xi, k)}{\sqrt{(a^2 - k^2)(1 - k^2)}} \quad (16)$$

where

$$\xi = \sin^{-1} \sqrt{\frac{a^2 - k^2}{a^2(1 - k^2)}} \quad (17)$$

In the ranges

$$k < a^2 < 1, \quad \theta < \xi$$

and when

$$k^2 < a^2 < k, \quad \theta > \xi.$$

In the program, Eq. (14) is used for  $\theta < \xi$  and Eq. (16) is used for  $\theta > \xi$ .

Case III:  $0 < a^2 < k^2$

$$\Pi(a^2, k) = K + \frac{a K Z(\beta, k)}{\sqrt{(1 - a^2)(k^2 - a^2)}} \quad (18)$$

where

$$\beta = \sin^{-1} \frac{a}{k} \quad (19)$$

Case IV:  $1 < a^2 < \infty$

$$\Pi(a^2, k) = - \frac{a K Z(A, k)}{\sqrt{(a^2 - 1)(a^2 - k^2)}} \quad (20)$$

where

$$A = \sin^{-1} \frac{1}{a} \quad (21)$$

The flow chart and the subroutine programmed in Fortran IV language\* are presented in Appendix C.

---

\* All programs in this report were written for the IBM 7090 Computer at Booth Computing Center, California Institute of Technology.

### 3. Incomplete Elliptic Integrals of the Third Kind

The incomplete elliptic integral of the third kind

$$\begin{aligned}\Pi(\varphi, a^2, k) &= \int_0^\varphi \frac{1}{1 - a^2 \sin^2 \theta} \frac{d\theta}{\sqrt{1 - k^2 \sin^2 \theta}} \\ &= \int_0^y \frac{1}{1 - a^2 t^2} \frac{dt}{\sqrt{(1 - t^2)(1 - k^2 t^2)}}\end{aligned}\quad (22)$$

has one more parameter than the complete one and consequently is more complicated. In general,  $\Pi(\varphi, a^2, k)$  can be classified as circular or hyperbolic, in the same way as the complete integral, however, in our subroutine, a different and more straightforward approach is taken.

First, we shall consider the special cases which can be integrated in terms of elementary functions and  $F(\varphi, k)$  and  $E(\varphi, k)$ .

$$\text{a) } \varphi = \frac{\pi}{2} ; \quad \Pi\left(\frac{\pi}{2}, a^2, k\right) = \Pi(a^2, k) \quad (23)$$

$$\text{b) } a^2 = 0 ; \quad \Pi(\varphi, 0, k) = F(\varphi, k) \quad (24)$$

$$\text{c) } k^2 = 0 ; \quad \left\{ \begin{array}{ll} \varphi & a^2 = 0 \end{array} \right. \quad (25a)$$

$$\left\{ \begin{array}{ll} \frac{1}{\sqrt{1 - a^2}} \tan^{-1} \left[ \sqrt{1 - a^2} \tan \varphi \right] & a^2 < 1 \end{array} \right. \quad (25b)$$

$$\left\{ \begin{array}{ll} \frac{1}{2\sqrt{a^2 - 1}} \log \left| \frac{\sqrt{a^2 - 1} \tan \varphi + 1}{\sqrt{a^2 - 1} \tan \varphi - 1} \right| & a^2 > 1 \end{array} \right. \quad (25c)$$

$$\left\{ \begin{array}{ll} \tan \varphi & a^2 = 1 \end{array} \right. \quad (25d)$$

$$d) \quad k = 1; \quad \left\{ \begin{array}{l} \frac{\sin \varphi}{2 \cos \varphi} + \frac{1}{4} \log \left[ \frac{1 + \sin \varphi}{1 - \sin \varphi} \right] \end{array} \right. \quad \alpha^2 = 1 \quad (26a)$$

$$\Pi(\varphi, \alpha^2, 1) = \left\{ \begin{array}{l} \frac{1}{1 - \alpha^2} \left[ \log(\tan \varphi + \sec \varphi) \right. \\ \left. - \frac{\alpha}{2} \log \left| \frac{1 + \alpha \sin \varphi}{1 - \alpha \sin \varphi} \right| \right] \end{array} \right. \quad \begin{array}{l} \alpha^2 > 0, \\ \alpha^2 \neq 1 \end{array} \quad (26b)$$

$$\left\{ \begin{array}{l} \frac{1}{1 - \alpha^2} \left[ \log(\tan \varphi + \sec \varphi) \right. \\ \left. + \sqrt{|\alpha^2|} \tan^{-1}(\sqrt{|\alpha^2|} \sin \varphi) \right] \end{array} \right. \quad \alpha^2 < 0 \quad (26c)$$

$$e) \quad \alpha^2 = 1; \quad \Pi(\varphi, 1, k) = \left[ k'^2 F(\varphi, k) - E(\varphi, k) + \tan \varphi \sqrt{1 - k^2 \sin^2 \varphi} \right] k'^{-2} \quad (27)$$

$$f) \quad \alpha^2 = k^2; \quad \Pi(\varphi, k^2, k) = \left[ E(\varphi, k) - \frac{k^2 \sin \varphi \cos \varphi}{\sqrt{1 - k^2 \sin^2 \varphi}} \right] k'^{-2} \quad (28)$$

The remaining values of  $\alpha^2$ , for  $0 < k^2 < 1$ , and  $0 < \varphi < \pi/2$  can be divided into three regions, with two series expansions [5]. In the regions

$$|\alpha^2| > 1, \quad k^2 < 1 \quad \text{and} \quad |\alpha^2| < 1, \quad k^2 < |\alpha^2|$$

the series expansion is

$$\Pi(\varphi, \alpha^2, k) = \sum_{m=0}^{\infty} b_m k^{2m}, \quad (29)$$

where

$$b_m = \binom{-\frac{1}{2}}{m} (-1)^m \int_0^\varphi \frac{\sin^{2m} \theta d\theta}{1 - \alpha^2 \sin^2 \theta}.$$

Hence

$$b_o = \begin{cases} \frac{1}{\sqrt{1-a^2}} \tan^{-1}[\sqrt{1-a^2} \tan \varphi] & a^2 < 1 \\ \frac{1}{2\sqrt{a^2-1}} \log \left| \frac{\sqrt{a^2-1} \tan \varphi + 1}{\sqrt{a^2-1} \tan \varphi - 1} \right| & a^2 > 1 \end{cases} \quad (30)$$

$$b_1 = \frac{b_o - \varphi}{2a^2}, \quad b_2 = \frac{1}{16a^4} [3a^2 \sin \varphi \cos \varphi + 6b_o - 3(2+a^2)\varphi];$$

and the recurrence relation is

$$2(m+1)a^2 b_{m+1} = (2m+1+2ma^2)b_m + (1-2m)b_{m-1} - (-)^m \binom{-\frac{1}{2}}{m-1} \sin^{2m-1} \varphi \cos \varphi \\ - (-)^m \binom{\frac{1}{2}}{m} \int_0^\varphi \sin^{2m} \theta d\theta.$$

In the region  $|a^2| < 1, \quad k^2 < 1$

a different series expansion is valid,

$$\Pi(\varphi, a^2, k) = \sum_{m=0}^{\infty} \sum_{j=0}^m (a^2)^m \binom{-\frac{1}{2}}{j} \left( \frac{k^2}{-a^2} \right)^j \int_0^\varphi \sin^{2m} \theta d\theta. \quad (32)$$

Both series can be derived very easily. The series in Eq. (29) is obtained by expanding the quantity

$$(1 - k^2 \sin^2 \theta)^{-\frac{1}{2}} = \sum_{m=0}^{\infty} \binom{-\frac{1}{2}}{m} (-k^2)^m \sin^{2m} \theta$$

and the series in Eq. (32), by the above expansion times the geometric series expansion of

$$(1 - a^2 \sin^2 \theta)^{-1} = \sum_{j=0}^{\infty} (a^2 \sin^2 \theta)^j.$$

Term by term integrations are then carried out for both expansions.

The regions of validity of the two series, with their overlapping

region  $1 > |a^2| > k^2$  are shown in Fig. 2.




Figure 2

We have chosen to evaluate the integral in this common region by Eq. (29). For values of  $\varphi$  and  $k$  such that  $k^2 \sin^2 \varphi$  is near one, both series expansions for  $\Pi(\varphi, a^2, k)$  converge slowly and hence certain addition formulae [5]<sup>\*</sup> must be used to facilitate these calculations. These addition formulae are of the form

$$\Pi(\varphi, a^2, k) \pm \Pi(\beta, a^2, k) = \Pi(\theta, a^2, k) \mp Q \quad (33)$$

where

$$\theta = 2 \tan^{-1} \left[ \frac{\sin \varphi \sqrt{1 - k^2 \sin^2 \beta} \pm \sin \beta \sqrt{1 - k^2 \sin^2 \varphi}}{\cos \varphi + \cos \beta} \right] \quad (34)$$

or

$$\theta = \cos^{-1} \left[ \frac{\cos \varphi \cos \beta \mp \sin \varphi \sin \beta \sqrt{(1 - k^2 \sin^2 \varphi)(1 - k^2 \sin^2 \beta)}}{1 - k^2 \sin^2 \varphi \sin^2 \beta} \right]. \quad (35)$$

<sup>\*</sup>

Some errors of sign for these formulae were found in this reference.



We now let

$$Q_a = \sqrt{\frac{a^2}{(1-a^2)(a^2-k^2)}} \tan^{-1} \left[ \frac{\sin \varphi \sin \beta \sin \theta \sqrt{a^2(1-a^2)(a^2-k^2)}}{1-a^2 \sin^2 \theta \pm a^2 \sin \varphi \sin \beta \cos \theta \sqrt{1-k^2 \sin^2 \theta}} \right] \quad (36)$$

and

$$Q_b = \sqrt{\frac{a^2}{(a^2-1)(a^2-k^2)}} \tanh^{-1} \left[ \frac{\sin \varphi \sin \beta \sin \theta \sqrt{a^2(a^2-1)(a^2-k^2)}}{1-a^2 \sin^2 \theta \pm a^2 \sin \varphi \sin \beta \cos \theta \sqrt{1-k^2 \sin^2 \theta}} \right]. \quad (37)$$

Then for

$$\begin{aligned} -\infty < a^2 < 0 & \quad , \quad Q = -Q_a; \\ 0 < a^2 < k^2 & \quad , \quad Q = Q_b; \\ k^2 < a^2 < 1 & \quad , \quad Q = Q_a; \\ 1 < a^2 < \infty & \quad , \quad Q = Q_b. \end{aligned} \quad (38)$$

In order to apply these formulae to our problem one must assign a value to  $\beta$ , choose a sign and then calculate a  $\theta$  from the given  $\varphi$  and  $k$ . The proper choice of  $\beta$  and sign will produce a  $\theta < \varphi$  and hence the series for  $\Pi(\theta, a^2, k)$  will converge with fewer terms than the series for  $\Pi(\varphi, a^2, k)$ .

Obviously the choice of  $\beta$  and the sign is rather important. The minimum  $\theta$  occurs when  $\beta = \pi/4$  and the lower sign is chosen. Making this substitution and choosing Eq. (34) the more convenient form for  $\theta$ , the addition formulae can be written as

$$\Pi(\varphi, a^2, k) = \Pi(\theta, a^2, k) + \Pi(\pi/4, a^2, k) + Q \quad (39)$$

where

$$\theta = 2 \tan^{-1} \left[ \frac{\sin \varphi \sqrt{1 - \frac{k^2}{2}} - \sqrt{\frac{1}{2}} (1 - k^2 \sin^2 \varphi)}{\cos \varphi + \sqrt{\frac{1}{2}}} \right] \quad (40)$$

and now

$$Q_a = \sqrt{\frac{a^2}{(1 - a^2)(a^2 - k^2)}} \tan^{-1} \left[ \frac{\sqrt{\frac{1}{2}} \sin \varphi \sin \theta \sqrt{a^2(1 - a^2)(a^2 - k^2)}}{1 - a^2 \sin^2 \theta - \frac{a^2}{\sqrt{2}} \sin \varphi \cos \theta \sqrt{1 - k^2 \sin^2 \theta}} \right], \quad (41)$$

$$Q_b = \sqrt{\frac{a^2}{(a^2 - 1)(a^2 - k^2)}} \tanh^{-1} \left[ \frac{\sqrt{\frac{1}{2}} \sin \varphi \sin \theta \sqrt{a^2(a^2 - 1)(a^2 - k^2)}}{1 - a^2 \sin^2 \theta - \frac{a^2}{\sqrt{2}} \sin \varphi \cos \theta \sqrt{1 - k^2 \sin^2 \theta}} \right], \quad (42)$$

and  $Q$  is given by Eq. (38).

This reduction in  $\theta$  is illustrated in Fig. 3 where  $\theta/\varphi$  is plotted as a function of  $\varphi$  for various values of  $k^2$ . At first glance one might decide that the addition formula should be applied whenever  $\varphi > \pi/4$ . It must be remembered, however, that whenever the addition formula is used the series for  $\Pi(\pi/4, a^2, k)$  must be evaluated as well as the one for  $\Pi(\theta, a^2, k)$  and hence the combined number of terms required must be compared against the number of terms required for the single series expansion of  $\Pi(\varphi, a^2, k)$ .




Figure 3. Variation of  $\theta/\phi$  with  $\phi$  for several values of the modulus  $k^2$ .  $\theta$  is calculated by Eq. (34).

For values of  $k^2$  very near one, the reduction in  $\varphi$  is not sufficient to assure fast convergence, and therefore repeated application of the addition formula can be carried out. Let  $\theta_n$  be the final value of  $\theta$  after the addition formula has been applied  $n$  times. Then the integral can be expressed as follows:

$$\Pi(\varphi, a^2, k) = n\Pi(\pi/4, a^2, k) + \Pi(\theta_n, a^2, k) + Q \quad (43)$$

where

$$\begin{aligned} \theta_1 &= 2 \tan^{-1} \left[ \frac{\sin \varphi \sqrt{1 - \frac{k^2}{2}} - \sqrt{\frac{1}{2}} (1 - k^2 \sin^2 \varphi)}{\cos \varphi + \sqrt{\frac{1}{2}}} \right], \\ \theta_2 &= 2 \tan^{-1} \left[ \frac{\sin \theta_1 \sqrt{1 - \frac{k^2}{2}} - \sqrt{\frac{1}{2}} (1 - k^2 \sin^2 \theta_1)}{\cos \theta_1 + \sqrt{\frac{1}{2}}} \right], \\ &\vdots \\ \theta_n &= 2 \tan^{-1} \left[ \frac{\sin \theta_{n-1} \sqrt{1 - \frac{k^2}{2}} - \sqrt{\frac{1}{2}} (1 - k^2 \sin^2 \theta_{n-1})}{\cos \theta_{n-1} + \sqrt{\frac{1}{2}}} \right]. \end{aligned} \quad (44)$$

so that,

$$\begin{aligned} Q_a &= \sqrt{\frac{a^2}{(1-a^2)(a^2-k^2)}} \left\{ \tan^{-1} \left[ \frac{\sqrt{\frac{1}{2}} \sin \varphi \sin \theta_1 \sqrt{a^2(1-a^2)(a^2-k^2)}}{1-a^2(\sin^2 \theta_1 + \sqrt{\frac{1}{2}} \sin \varphi \cos \theta_1 \sqrt{1-k^2 \sin^2 \theta_1})} \right] \right. \\ &\quad + \dots + \dots \\ &\quad \left. + \tan^{-1} \left[ \frac{\sqrt{\frac{1}{2}} \sin \varphi \sin \theta_n \sqrt{a^2(1-a^2)(a^2-k^2)}}{1-a^2(\sin^2 \theta_n + \sqrt{\frac{1}{2}} \sin \varphi \cos \theta_n \sqrt{1-k^2 \sin^2 \theta_n})} \right] \right\}, \end{aligned} \quad (45)$$

and

$$Q_b = \sqrt{\frac{a^2}{(a^2 - 1)(a^2 - k^2)}} \left\{ \tanh^{-1} \left[ \frac{\sqrt{\frac{1}{2}} \sin \varphi \sin \theta_1 \sqrt{a^2(a^2 - 1)(a^2 - k^2)}}{1 - a^2(\sin^2 \theta_1 + \sqrt{\frac{1}{2}} \sin \varphi \cos \theta_1 \sqrt{1 - k^2 \sin^2 \theta_1})} \right] \right. \\
+ \dots + \dots + \dots + \left. \tanh^{-1} \left[ \frac{\sqrt{\frac{1}{2}} \sin \varphi \sin \theta_n \sqrt{a^2(a^2 - 1)(a^2 - k^2)}}{1 - a^2(\sin^2 \theta_n + \sqrt{\frac{1}{2}} \sin \varphi \cos \theta_n \sqrt{1 - k^2 \sin^2 \theta_n})} \right] \right\}. \quad (46)$$

Again  $Q$  is given by Eq. (38).

Table I illustrates how well repeated application of the addition formula works for the most difficult cases expected to be encountered. In this table values of  $\theta_n$  for  $n = 1, 2, 3, 4, 5$  are shown for  $\varphi = \pi/2$  and  $k^2 = 0.9990$  and  $0.9999$ .

Table I Reduction of the Argument

$k^2$	$\varphi$	$\theta_1$	$\theta_2$	$\theta_3$	$\theta_4$	$\theta_5$
0.9990	$\pi/2$	1.5392	1.4820	1.3518	1.0499	0.4271
0.9999	$\pi/2$	1.5608	1.5425	1.5008	1.4015	1.1664

Test runs were made with our program and the total number of terms were recorded for cases when the addition formula was used and when it was not used. We found that the least computer time resulted if we used the addition formula for values of  $\varphi$  and  $k$  such that

$k \sin \varphi > 0.75$ . Furthermore, the addition formula is repeatedly used until  $k \sin \theta_n < 0.75$ . In this way we were able to calculate  $\Pi(\varphi, a^2, k)$  for all ranges of  $a^2$ ,  $|a^2 - k^2| > 0.1$ ,<sup>\*</sup> where  $0 < \varphi < 1.570$  and  $0 < k^2 < 0.999$  using less than a total of 25 terms.

The flow chart and a copy of the subroutine are given in Appendix C.

---

<sup>\*</sup> The details of the region  $|a^2 - k^2| \leq 0.1$  are given in Appendix C.

## APPENDICES

Appendix A. Approximate functions for the complete elliptic integrals of the first and second kind.

The complete elliptic integrals of the first and second kind, i. e. ,

$$K(k) = \int_0^{\pi/2} \frac{d\theta}{\sqrt{1 - k^2 \sin^2 \theta}} \quad (A.1)$$

$$E(k) = \int_0^{\pi/2} \sqrt{1 - k^2 \sin^2 \theta} \, d\theta \quad (A.2)$$

where  $0 \leq k^2 < 1$ , can be calculated by the approximate functions [3]

$$\begin{aligned} K^*(k) = & \{a_0 + a_1 \eta + a_2 \eta^2 + a_3 \eta^3 + a_4 \eta^4\} \\ & + \{b_0 + b_1 \eta + b_2 \eta^2 + b_3 \eta^3 + b_4 \eta^4\} \log \frac{1}{\eta} \end{aligned} \quad (A.3)$$

and

$$\begin{aligned} E^*(k) = & \{1 + c_1 \eta + c_2 \eta^2 + c_3 \eta^3 + c_4 \eta^4\} \\ & + \{d_1 \eta + d_2 \eta^2 + d_3 \eta^3 + d_4 \eta^4\} \log \frac{1}{\eta} \end{aligned} \quad (A.4)$$

In these equations

$$\eta = 1 - k^2 = k'^2$$

and the coefficients as given in Ref. [3] are

$$a_0 = 1.3862,9436,112$$

$$a_1 = 0.0966,6344,259$$

$$a_2 = 0.0359,0092,383$$

$$a_3 = 0.0374,2563,713$$

$$a_4 = 0.0145,1196,212$$

$$c_1 = 0.4432,5141,463$$

$$c_2 = 0.0626,0601,220$$

$$c_3 = 0.0475,7383,546$$

$$c_4 = 0.0173,6506,451$$

$$b_0 = 0.5$$

$$b_1 = 0.1249,8593,597$$

$$b_2 = 0.0688,0248,576$$

$$b_3 = 0.0332,8355,346$$

$$b_4 = 0.0044,1787,012$$

$$d_1 = 0.2499,8368,310$$

$$d_2 = 0.0920,0180,037$$

$$d_3 = 0.0406,9697,526$$

$$d_4 = 0.0052,6449,639$$

The error curves of these approximation functions are shown in Fig. A-1.




Figure A-1. Error in  $K^*$  and  $E^*$  as a function of  $\sin^{-1}k$ . This figure is given in Ref. [3], p 172.



The subroutines of these functions are not included since they are quite straightforward.

## Appendix B: The Evaluation of the Incomplete Elliptic Integrals of the First and Second Kind.

The incomplete elliptic integrals of the first and second kind

$$F(\varphi, k) = \int_0^{\varphi} \frac{d\theta}{\sqrt{1 - k^2 \sin^2 \theta}} \quad (\text{B.1a})$$

$$E(\varphi, k) = \int_0^{\varphi} \sqrt{1 - k^2 \sin^2 \theta} \, d\theta \quad (\text{B.1b})$$

are evaluated by two types of series and some special formulae for the boundary condition cases. The reader may refer to Ref. 4 for detailed discussion of these series expansions.

In the range  $0 < |k \sin \varphi| < \tanh 1 = 0.7615, 942$  the series expansions are evaluated by the equations

$$F(\varphi, k) = \sum_{n=0}^{\infty} \frac{(2n)!}{2^{2n} n!^2} k^{2n} \int_0^{\varphi} \sin^{2n} \theta \, d\theta \quad (\text{B.2a})$$

and

$$E(\varphi, k) = - \sum_{n=0}^{\infty} \frac{(2n)!}{2^{2n} n!^2 (2n-1)} k^{2n} \int_0^{\varphi} \sin^{2n} \theta \, d\theta \quad (\text{B.2b})$$

These formulae can be derived directly by the binomial expansion of the integrands of  $F(\varphi, k)$  and  $E(\varphi, k)$  in Eqs. (B.1). Though these series are convergent for the entire range of  $\varphi$  and  $k$ , their convergence is

slow when  $|k \sin \varphi| > \tanh 1$ . A second set of series therefore is used to evaluate  $F(\varphi, k)$  and  $E(\varphi, k)$  in the range

$$\tanh 1 < |k \sin \varphi| < 1.$$

These are presented in Ref. 4 and are

$$\begin{aligned} F(\varphi, k) = & \frac{2}{\pi} K' \log \left[ \frac{4}{\sqrt{1-k^2 x^2} + |k| \sqrt{1-x^2}} \right] + |k| \sqrt{\frac{1-x^2}{1-k^2 x^2}} \log \left[ \frac{1+|kx|}{2} \right] \\ & - |k| \sqrt{(1-x^2)(1-k^2 x^2)} \sum_{n=0}^{\infty} \frac{2^{2n} n!^2}{(2n+1)!} (1-k^2 x^2)^n \sum_{m=n+1}^{\infty} \left[ \frac{(2m+2)!}{2^{2m+2} (m+1)!^2} \right]^2 (k')^{2m-2n} \\ & - \sum_{n=1}^{\infty} \left[ \frac{(2n)!}{2^{2n} n!^2} \right]^2 (k')^{2n} \sum_{m=1}^n \frac{1}{m(2m-1)}, \end{aligned} \quad (\text{B.3a})$$

$$\begin{aligned} E(\varphi, k) = & \frac{2}{\pi} (K' - E') \log \left[ \frac{4}{\sqrt{1-k^2 x^2} + |k| \sqrt{1-x^2}} \right] - |k| \sqrt{\frac{1-x^2}{1-k^2 x^2}} (1 - |kx|) \\ & - |k| \sqrt{(1-x^2)(1-k^2 x^2)} \sum_{n=0}^{\infty} \frac{2^{2n} n!^2}{(2n+1)!} (1-k^2 x^2)^n \sum_{m=n+1}^{\infty} \left[ \frac{(2m)!}{2^{2m} m!^2} \right]^2 \frac{2m+1}{2m+2} (k')^{2m-2n} \\ & - \sum_{n=1}^{\infty} \left[ \frac{(2n)!}{2^{2n} n!^2} \right]^2 \frac{2n}{(2n-1)} (k')^{2n} \sum_{m=1}^n \frac{1}{m(2m-1)} + \sum_{n=0}^{\infty} \left[ \frac{(2n)!}{2^{2n} n!^2} \right]^2 \frac{1}{(2n-1)^2} (k')^{2n} \end{aligned} \quad (\text{B.3b})$$

where

$$x = \sin \varphi.$$

Our subroutine for  $F(\varphi, k)$  and  $E(\varphi, k)$  follows closely the one suggested in Ref. 4, pages 11 to 14. The flow chart and a copy of the subroutine written in Fortran IV language for the IBM 7090 are given in Appendix C.

The maximum number of terms required to yield six decimal places in the neighborhood of  $k \sin \varphi = \tanh 1$  is nineteen terms for the first series method (Eqs. (B.2)), and fourteen terms for the second method (Eqs. (B.3)). It should also be pointed out that even though Eqs. (B.3) are theoretically good in the regions of  $k \sin \varphi = 1$ , when  $k^2 \sin^2 \varphi$  exceeds 0.99 the number of significant figures is reduced. This loss in accuracy occurs because the quantity  $1 - k^2 \sin^2 \varphi$  must be evaluated.

A few words are also needed for the boundary condition cases:

Case 1:  $k^2$  near 0

This presents no special problem since the first series (B.2) will converge rapidly. We have arbitrarily considered  $k^2 < 10^{-7}$  to be equal to zero and hence only the first terms of the series in Eqs. (B.2) are required yielding

$$F(\varphi, 0) = \varphi$$

and

$$E(\varphi, 0) = \varphi \quad .$$




Figure C-1. The flow chart of Subroutine PX for calculation of the complete or incomplete elliptic integral of the third kind. The circled numbers in the flow chart refer to statement numbers in the program listing in Fig. C-2.

Case 2:  $k^2 \sim 1$  or  $k'^2 = 1 - k^2 \sim 0$

$F(\varphi, k)$  and  $E(\varphi, k)$  have the following series expansions [5]  
in powers of  $k'^2$

$$F(\varphi, k) = \sum_{m=0}^{\infty} \left( \frac{-\frac{1}{2}}{m} \right) k'^{2m} \rho_{2m}(\varphi) \quad [0 < k'^2 \tan^2 \varphi < 1, k^2 < 1] \quad (\text{B.4a})$$

where

$$\rho_0(\varphi) = \log \frac{1 + \sin \varphi}{\cos \varphi}$$

$$\rho_2(\varphi) = \frac{\sin \varphi \sec^2 \varphi}{2} - \frac{1}{2} \log \frac{1 + \sin \varphi}{\cos \varphi}$$

$$\rho_4(\varphi) = \frac{1}{8} [2 \sin^3 \varphi \sec^4 \varphi - 3 \sin \varphi \sec^2 \varphi + 3 \log \frac{1 + \sin \varphi}{\cos \varphi}]$$

.

$$\rho_{2m}(\varphi) = \frac{1}{2m} [\sin^{2m-1} \varphi \sec^{2m} \varphi + (1 - 2m) \rho_{2m-2}(\varphi)]$$

and

$$E(\varphi, k) = \sum_{m=0}^{\infty} \left( \frac{\frac{1}{2}}{m} \right) k'^{2m} d_{2m}(\varphi) \quad [0 < k'^2 \tan^2 \varphi < 1; k^2 < 1] \quad (\text{B.4b})$$

where

$$d_0(\varphi) = \sin \varphi$$

$$d_2(\varphi) = -\sin \varphi + \log \frac{1 + \sin \varphi}{\cos \varphi}$$

$$d_4(\varphi) = \frac{1}{2} [\sin^3 \varphi \sec^2 \varphi + 3 \sin \varphi - 3 \log \frac{1 + \sin \varphi}{\cos \varphi}]$$

.

$$d_{2m}(\varphi) = \frac{1}{2(m-1)} [\sin^{2m-1} \varphi \sec^{2(m-1)} \varphi + (1 - 2m) d_{2(m-1)}(\varphi)]$$

with  $m \neq 1$ .

Again we have arbitrarily considered  $10^{-7}$  our criterion for zero. That is we have taken  $|k'^2| \leq 10^{-7}$  to be  $k'^2 = 0$  or  $|1.0 - k^2| \leq 10^{-7}$  to be the same as  $k^2 = 1.0$  so that  $F$  and  $E$  are represented by their first terms in Eqs. (B.4) giving

$$F(\varphi, 1) = \log (\tan \varphi + \sec \varphi)$$

and

$$E(\varphi, 1) = \sin \varphi .$$

The flow chart and Fortran IV program are given in Appendix C.

Appendix C. The most frequently used symbols in the subroutine listings in Figs. C-2 and C-4 have the following definitions:

AS	$a^2$
XK	$k$
SK	$k^2$
PK	$k'$
PHI	$\varphi$
THE	$\theta$
K	$K^*$
Ec	$E^*$
F	$F(\varphi, k)$
E	$E(\varphi, k)$
PI	$\Pi(\varphi, a^2, k)$ or $\Pi(a^2, k)$
TOL	Allowable error
M	Accuracy or error indicator - (output)  M = 1, less than four figure accuracy or error return due to non-convergence, overflow or illegal parameter in argument  M = 2, six figure accuracy  M = 3, four to six figure accuracy
I	a) Indicator in subroutine PIX - (output)  I = 0, $\theta$ , F, and E were not calculated  I = 1, $\theta$ , $F(\theta, k')$ and $E(\theta, k')$ were calculated  I = 2, $\theta$ , $F(\theta, k)$ and $E(\theta, k)$ were calculated

b) Indicator in subroutine ELLI - (input)

I = 1, both F and E to be calculated

I = 2, calculate E only

I = 3, calculate F only

Figures C-1 and C-2 are the flow chart and Fortran IV listing for the subroutine which calculates  $\Pi(a^2, k)$  or  $\Pi(\varphi, a^2, k)$  the complete or incomplete elliptic integral of the third kind. The circled numbers in Fig. C-1 refer to statement numbers in the program listing given in Fig. C-2.

Six figure accuracy is guaranteed except when  $|1 - a^2| < 0.1$  or  $k^2 > 0.9$  and at least four figure accuracy is guaranteed when  $|1 - a \sin \varphi| \geq 0.0001$  and  $k \sin \varphi \leq 0.9999$ . The accuracy indicator M is set as described above and a message is printed whenever the accuracy is less than six figures.

If  $k \sin \varphi$  is greater than 0.75 or  $\varphi = \pi/2$  the methods outlined in the text cannot calculate  $\Pi(a^2, k)$  or  $\Pi(\varphi, a^2, k)$  to six figures when  $|a^2 - k^2| \leq 0.1$  or to four figures when  $|a^2 - k^2| \leq 0.0001$ , because accuracy is lost in calculating the quantity  $a^2 - k^2$ . Therefore, our program transfers control to statement number 101 using Eq. (32), without the addition formulae for both integrals whenever  $|a^2 - k^2|$  is less than or equal to 0.1 and  $M = 2$  (six figure accuracy); thus the quantity  $a^2 - k^2$  need not be evaluated. When  $|a^2 - k^2| \leq 0.0001$  and no more than four figure accuracy is required ( $M = 1$  or  $M = 3$ ),  $a^2$  is considered to be the same as  $k^2$  so that

$$\Pi(\varphi, a^2, k) = \Pi(\varphi, k^2, k)$$



or

$$\Pi(a^2, k) = \Pi(k^2, k).$$

The remaining deviation from the text is the case when  $\varphi = \pi/2$ ,  $M = 3$  and  $0.0001 < |a^2 - k^2| < 0.1$ . For this case better accuracy is obtained if  $\Pi(a^2, k)$  is calculated as a special case of the incomplete integral.

The flow chart and Fortran IV listing for the subroutine which calculates the incomplete elliptic integrals of the first and second kind are given in Figs. C-3 and C-4 respectively. As above, the circled numbers in the flow chart correspond to statement numbers in the program listing.

Figure C-2. The Fortran IV listing of Subroutine PIX.

```

SUBROUTINE PIX(PHI,AS,XK,XL,TOL,M,PI,K,EG,I,THE,F,E)
  1  FORMAT(19H PH, K GT 1, NEG OR GT PI/2)
  2  FORMAT(7H K GT 1, NEG OR GT PI/2)
  3  FORMAT(130H PI AT LEAST 4 FIGURE ACCURACY)
  4  FORMAT(150H OVERFLOW IN PI)
  5  FORMAT(17H PI NOT CONVERGED)
  6  FORMAT(120H PI NEAR SINGULARITY)
  7  FORMAT(122H AS OR SK=ONE, PIC=INF)
1007  REAL K,N
      DIMENSION B(101)
      TAL=0.
      PIT=0.
      Q=0.
      NN=1
      THE=0.
      PI=0.
      N=0.
      EC=0.
      K=0.
      F=0.
      E=0.
      CALL OVERFL(L)
      IF (ABS(PHI).LE..0000001) GO TO 10
      SP=K*XK
      ST=SIN(PHI)
      AS=K*AS
      ASK=ABS(AS)
      IF (AS-.01) ASP=ASP
      IF (ABS(1.0-AS)) GE .1 AND .SK.LE..9) GO TO 8
      IF (ABS(1.0-ASP).LT...0001.OR.XK*SP.GT..9999) GO TO 7
      WRITE(6,3)
      M=3
      GO TO 9
      7  WRITE(6,6)
      M=1
      GO TO 9
      8  M=2
      IF (.0000001.LT.PHI.AND.PHI.LT.1.5707962) GO TO 20
      IF (ABS(PHI-1.5707963).LE..0000001) GO TO 1000
      WRITE(6,1)
      M=1
      RETURN
      10  PI=0.
      N=2
      RETURN
      20  CP=COS(PHI)
      SPS=SP*SP
      IF (.0000001.LT.SK.AND.SK.LE..9999) GO TO 30
      IF (SK.LE..0000001) GO TO 40
      IF (SK.LT.1.0000001) GO TO 50
      WRITE (6,2)
      M=1
      RETURN
      40  IF (ABS(AS).LE..0000001) GO TO 43
      IF (ABS(AS-1.0).LT..0001) GO TO 41
      X=SQRT(1.0) GO TO 42
      PI=ATAN(X*SP/CP)/X
      RETURN
      41  PI=SP/CP
      RETURN
      42  X=SQRT(AS-1.0)
      PI=.5/X*ALOG(ABS(((X*SP/CP)+1.)/(X*SP/CP)-1.0)))
      RETURN
      43  PI=PI
      M=2
      RETURN
      50  IF (ABS(AS-1.0).LT..0001) GO TO 52
      IF (AS.LT..0.) GO TO 53
      PI=1/ALOG((SP+1.0)/CP)-0.5*A*LOG(ABS((1.0+ASP)/(1.0-ASP)))
      1  (1.0-AS)
      RETURN
      52  PI=.5*(SP/(CP*CP)+.5*A*LOG((1.0+SP)/(1.0-SP)))
      RETURN
      53  PI=1/ALOG((SP+1.0)/CP)+A*ATAN(-ASP)/(1.0-AS)
      RETURN
      30  IF (ABS(AS).LE..0000001) GO TO 31

```

Figure C-2 continued

```

C=-.5*AK
CM=1.0*C
PI=CM*AT*T
DO 110 MM=2,200
N=MM
TN=2*MM
SC=SC*SPS
T=((TN-1.0)*T-SC)/TN
AT=AT*AS
C=C*(10.5-N)*AK/N
SM=CM*AT
PI=PI*SM
IF(ABS(SW/(PI*PHI))-LE,TOL) GO TO 103
IF(ABS(1/T)-GT,1.0 E-30) GO TO 110
AT=AT*1.0 E30
C=C*1.0 E-30
CM=CM*1.0 E-30
110 CONTINUE
GO TO 261
103 PI=PI*PHI
GO TO (501,121,122),NN
102 THE=2.0*ATAN((SQRT(1.0--5*SK)*SP-.70710678*SQRT(1.0-SK*SPS))/
1 (CP+.70710678))
ST=SIN(THE)
CT=COS(THE)
ST=ST*ST
CT=CT*CT
Y=(AS*ST-TOL) GO TO 105
Y=-.70710678*SP*ST*SQRT(AS*(AS-1.0)*(AS-SK))/
1 (1.0-AS*STS-.70710678*SP*CT*SQRT(1.0-SK*STS)*AS)
TY=-5*ALOG(ABS((1.0*Y)/(1.0-Y)))
Q=SQRT(AS/((AS-1.0)*(AS-SK)))*TY+Q
104 PHI=THE
SP=ST
CP=CT
SPS=STS
TAL=TAL+1.0
IF(XX*SP-GT,.75) GO TO 102
NN=2
GO TO 101
105 Q=-SQRT(AS/((1.0-AS)*(AS-SK)))*ATAN(.70710678*SP*ST*SQRT(AS*(1.0-
1 AS)*(AS-SK))/((1.0-AS*STS-.70710678*AS*SP*CT*SQRT(1.0-SK*STS)))+Q
GO TO 104
121 PI=PI
PHI=78539816
SP=.70710678
CP=-.70710678
SPS=-.5
GO TO 101
122 PI=TAL*PI+PI+Q
RETURN
300 IF(XX*SP-GT,.75) GO TO 305
302 SAD=SQRT(AS-1.0)
XX=SAD*SP/CP
BO=-.5/SAD*ALOG(ABS((XX+1.0)/(XX-1.0)))
GO TO 251
305 THE=2.0*ATAN((SQRT(1.0--5*SK)*SP-.70710678*SQRT(1.0-SK*SPS))/
1 (CP+.70710678))
ST=SIN(THE)
CT=COS(THE)
ST=ST*ST
CT=CT*CT
Y=.70710678*SP*ST*SQRT(AS*(AS-1.0)*(AS-SK))/
1 (1.0-AS*STS-.70710678*SP*CT*SQRT(1.0-SK*STS)*AS)
TY=-5*ALOG(ABS((1.0*Y)/(1.0-Y)))
Q=SQRT(AS/((AS-1.0)*(AS-SK)))*TY+Q
PHI=THE
SP=ST
CP=CT
SPS=STS
TAL=TAL+1.0
IF(XX*SP-GT,.75) GO TO 305
NN=4
GO TO 302
310 PI=PI
NN=5
PHI=.78539816
SP=.70710678

```

```

CP=-.70710678
SPS=-.5
GO TO 302
311 PI=TAL*PI+PI+Q
501 CALL OVERFL(L)
IF(L.EQ.2) RETURN
WRITE(6,4)
M=1
RETURN
1000 IF(ABS(1-1.0)*LE-.0000001.OR,ABS(SK-1.0)*LE-.0000001) GO TO 1050
IF(1.0000001.LT,SK.AND,SK.LT.99999999) GO TO 1025
WRITE(6,4)
1006 IF(1.0000001.LT,SK) GO TO 1040
M=1
RETURN
1050 WRITE(6,1007)
M=1
RETURN
1040 IF(AS.LT.-99999999) GO TO 1041
PI=0.
M=2
RETURN
1041 PI=1.5707963/SQRT(1.0-AS)
RETURN
1025 PK=SQRT(1.0-SK)
CALL ELLPT(XK,KECL)
IF(ABS(1AS*(AS-XK)*LE-.0000001) GO TO 1031
IF(ABS(1AS*(AS-XK)*LE-.0000001) GO TO 1032
IF(ABS(1AS*(AS-XK)*LE-.0000001) GO TO 1033
IF(ABS(1AS*(AS-XK)*LE-.0000001) GO TO 1033
SPS=1.0
CP=0.
IF(ABS(AS-SK)*LE-.0001) GO TO (1033,101,1033),M
1026 IF(AS-GT,SK.AND,AS-LE-.9999) GO TO 1200
IF(AS-LT,SK.AND,AS-GT-.0000001) GO TO 1300
THE=ASIN(SQRT(1.0/AS))
CALL ELLI(THE,XX,F,E,1,L,TOL)
PI=(EC*F-K*E)*SQRT(AS/((AS-1.0)*(AS-SK)))
I=2
RETURN
1031 PI=K
RETURN
1032 PI=.0.25*(3.14159265+2.0*(1.0-AS)*K)/(1.0-AS)
RETURN
1033 RETURN
1100 XK=-XK
IF(AS-LT,XXK) GO TO 1102
THE=ASIN(SQRT(AS/AS*(AS-SK)))
CALL ELLI(THE,PK,F,E,1,L,TOL)
PI=SK*K/(SK-AS)-AS*(EC*F+K*(E-F))/SQRT(AS*(1.0-AS)*(AS-SK))
I=1
RETURN
1102 THE=ASIN(1.0/SQRT(1.0-AS))
CALL ELLI(THE,PK,F,E,1,L,TOL)
PI=K/(1.0-AS)+AS*(EC*F+K*(E-F)-1.5707963)/
1 SQRT(AS*(1.0-AS)*(AS-SK))
I=1
RETURN
1200 IF(AS-GT,XXK) GO TO 1203
IF(AS-LT,XXK) GO TO (1203,101,1203),M
CALL ELLI(THE,PK,F,E,1,L,TOL)
PI=(EC*F+K*(E-F))*SQRT(AS/((AS-SK)*(1.0-AS)))
I=1
RETURN
1203 THE=ASIN(SQRT((1.0-AS)/(1.0-SK)))
CALL ELLI(THE,PK,F,E,1,L,TOL)
PI=K+(1.5707963-(EC*F+K*(E-F)))*SQRT(AS/((AS-SK)*(1.0-AS)))
I=1
RETURN
1300 THE=ASIN(SQRT(AS/AS*(AS-SK)))
CALL ELLI(THE,XX,F,E,1,L,TOL)
PI=K*(K*E-EC*F)*SQRT(AS/((1.0-AS)*(SK-AS)))
I=2
RETURN
END

```

For Figures C -3 and C -4 the captions should be interchanged.

Figure C-3. The flow chart of Subroutine ELLI for calculation of the incomplete elliptic integrals of the first and second kinds. The circled numbers in the flow chart refer to statement numbers in the program listing in Fig. C-4.

```

SUBROUTINE ELLI(PHI,XK,F,E,I,M,TOL)
REAL N
CALL OVERFL(M)
F=0.
E=0.
SK=XK*XK
IF(1.0000001.LT.SK.AND.SK.LT..99999999) GO TO 25
IF(SK.LE..0000001) GO TO 20
IF(ABS(XK-1.0).LE..0000001) GO TO 23
19 WRITE(6,3)
3 FORMAT(7H K GT 1)
M=1
RETURN
20 E=PHI
F=PHI
M=2
RETURN
23 SP=SIN(PHI)
E=SP
M=2
IF(1.EQ.2) RETURN
F=ALOG(SP/(COS(PHI))+1.0/(COS(PHI)))
10 CALL OVERFL(M)
IF(M.EQ.1) WRITE(6,4)
4 FORMAT(17H OVERFLOW IN ELLI)
RETURN
25 IF(.0000001.LT.PHI.AND.PHI.LT.1.5707960) GO TO 46
IF(ABS(PHI).LE..0000001) GO TO 41
IF(ABS(PHI-1.5707963).LE..0000003) GO TO 44
40 WRITE(6,5)
5 FORMAT(23H NEG PHI OR PHI GT PI/2)
M=1
RETURN
41 E=0.
F=0.
M=2
RETURN
44 CALL ELIPT(XK,F,E,M)
RETURN
46 SP=SIN(PHI)
AKP=XK*SP
IF(AKP.GT..7615942) GO TO 45
43 SK=XK*XK
CE=-.5*SK
CF=-CE
SPS=SP*SP
CP=COS(PHI)
SCT=SP*CP
T=.5*(PHI-SCT)
E=CE*T
F=CF*T
DO 64 J=2,200
N=J
TN=2.0*N
SCT=SCT*SPS
T=((TN-1.0)*T-SCT)/TN
A=-SK/N
CF=CF*(1.5-N)*A
CE=CE*(1.5-N)*A
TCE=CE*T
TCF=CF*T
F=F+TCF
GO TO (62,61,62),I
62 IF(ABS(TCF).LE.TOL) GO TO 65
GO TO 64
61 IF(ABS(TCE).LE.TOL) GO TO 66
64 CONTINUE
M=1
WRITE(6,6)
6 FORMAT(19H ELLB NOT CONVERGED)
RETURN
65 F=PHI+F
66 E=PHI+E
GO TO 10
45 A=XK*COS(PHI)
B2=1.0-AKP*AKP
IF(B2.GE..05) GO TO 48
IF(B2.LT..0005) GO TO 47
WRITE(6,1)
1 FORMAT(32H F AND E AT LEAST 4 FIG.ACCURACY)
M=3
GO TO 49
47 WRITE(6,2)
2 FORMAT(34H F AND E LESS THAN 4 FIG. ACCURACY)
M=1
GO TO 49
48 M=2
49 B=SQRT(B2)
ABL=ALOG(4.0/(A+B))
ADB=A/B
PKS=1.0-XK*XK
X1=.5
XJ=.25*PKS
XL=1.0
AB=A*B
XM=-AB*PKS*.140625
XN=-AB*PKS*.1875
S1=XM-XJ*XL
S2=XN-.25*PKS
S3=XJ
S4=.75*XJ
DO 200 J=2,100
N=J
TN=2*J
C1=TN-1.0
C2=C1/TN
C3=(TN+1.0)/(TN+2.0)
C4=C3*PKS
AB=AB*B2
CN=AB*X1/TN
CD2A=C2*PKS*XJ
CD2B=PKS*XJ/(TN*TN)
X1=C2*X1
XJ=C3*C4*XJ
XL=XL+1.0/(N*C1)
XM=(XM-AB*X1/TN)*C3*C4
XN=(XN-CN)*C4*C2
D1=XM-XJ*XL
D2=XN-CD2A*XL+CD2B
S1=S1+D1
S2=S2+D2
S3=S3+XJ
S4=S4+C3*XJ
GO TO (90,91,90),I
90 IF(ABS(D1).LE.TOL) GO TO 210
GO TO 200
91 IF(ABS(D2).LE.TOL) GO TO 211
200 CONTINUE
M=1
WRITE(6,6)
RETURN
210 F=(1.0+S3)*ABL+ADB*ALOG(.5+.5*AKP)+S1
IF(1.EQ.3) RETURN
211 E=(.5+S4)*PKS*ABL+1.0-ADB*(1.0-AKP)+S2
RETURN
END

```




Figure C-4. The Fortran IV listing of Subroutine ELLI.

### REFERENCES

1. Selfridge, R. G., and Maxfield, J. E., "A Table of the Incomplete Elliptic Integral of the Third Kind", Dover, 1958.
2. Belyakov, V. M., Kravtsova, R. N., and Rappoport, M. G. "Tables of Elliptic Integrals", Vol. 1, Academy NAUK SSSR, Moscow, 1962.
3. Hastings, Jr., Cecil, "Approximations for Digital Computer", Princeton University Press, 1955.
4. DiDonato, A. R., and Hershey, A. V., "New Formulae for Computing Incomplete Elliptic Integrals of the First and Second Kind", Computation and Exterior Ballistics Laboratory, U. S. Naval Proving Ground, Dahlgren, Virginia, NAVORD Report No. 5906, January 1959.
5. Byrd, P. F., and Friedman, M. D., "Handbook of Elliptic Integrals for Engineers and Physicists", Springer-Verlag, Berlin, 1954.

# DISTRIBUTION LIST FOR UNCLASSIFIED TECHNICAL REPORTS

ISSUED UNDER

CONTRACT Nonr-220(43)

(Single copies unless otherwise specified)

Chief of Naval Research  
Department of the Navy  
Washington 25, D. C.  
Attn: Codes 438 (3)

461

463

466

Commanding Officer  
Office of Naval Research  
Branch Office  
495 Summer Street  
Boston 10, Massachusetts

Commanding Officer  
Office of Naval Research  
Branch Office  
207 West 24th Street  
New York 11, New York

Commanding Officer  
Office of Naval Research  
Branch Office  
1030 East Green Street  
Pasadena, California

Commanding Officer  
Office of Naval Research  
Branch Office  
1000 Geary Street  
San Francisco 9, California

Commanding Officer  
Office of Naval Research  
Branch Office  
Box 39, Navy No. 100  
Fleet Post Office  
New York, New York (25)

Director  
Naval Research Laboratory  
Washington 25, D. C.  
Attn: Code 2027 (6)

Chief, Bureau of Naval Weapons  
Department of the Navy  
Washington 25, D. C.  
Attn: Codes RUAW-r  
RRRE  
RAAD  
RAAD-222  
DIS-42

Commander  
U. S. Naval Ordnance Test Station  
China Lake, California  
Attn: Code 753

Chief, Bureau of Ships  
Department of the Navy  
Washington 25, D. C.

Attn: Codes 310

312

335

420

421

440

442

449

Chief, Bureau of Yards and Docks  
Department of the Navy  
Washington 25, D. C.  
Attn: Code D-400

Commanding Officer and Director  
David Taylor Model Basin  
Washington 7, D. C.

Attn: Codes 108

142

500

513

520

525

526

526A

530

533

580

585

589

591

591A

700

Commander  
U.S. Naval Ordnance Test Station  
Pasadena Annex  
3202 E. Foothill Blvd.  
Pasadena 8, California  
Attn: Code P-508

Commander  
Planning Department  
Portsmouth Naval Shipyard  
Portsmouth, New Hampshire

Commander  
Planning Department  
Boston Naval Shipyard  
Boston 29, Massachusetts



Commander  
Planning Department  
Pearl Harbor Naval Shipyard  
Navy No. 128, Fleet Post Office  
San Francisco, California

Commander  
Planning Department  
San Francisco Naval Shipyard  
San Francisco 24, California

Commander  
Planning Department  
Mare Island Naval Shipyard  
Vallejo, California

Commander  
Planning Department  
New York Naval Shipyard  
Brooklyn 1, New York

Commander  
Planning Department  
Puget Sound Naval Shipyard  
Bremerton, Washington

Commander  
Planning Department  
Philadelphia Naval Shipyard  
U. S. Naval Base  
Philadelphia 12, Pennsylvania

Commander  
Planning Department  
Norfolk Naval Shipyard  
Portsmouth, Virginia

Commander  
Planning Department  
Charleston Naval Shipyard  
U. S. Naval Base  
Charleston, South Carolina

Commander  
Planning Department  
Long Beach Naval Shipyard  
Long Beach 2, California

Commander  
Planning Department  
U. S. Naval Weapons Laboratory  
Dahlgren, Virginia

Commander  
U. S. Naval Ordnance Laboratory  
White Oak, Maryland

Dr. A. V. Hershey  
Computation and Exterior  
Ballistics Laboratory  
U. S. Naval Weapons Laboratory  
Dahlgren, Virginia

Superintendent  
U. S. Naval Academy  
Annapolis, Maryland  
Attn: Library

Superintendent  
U. S. Naval Postgraduate School  
Monterey, California

Commandant  
U. S. Coast Guard  
1300 E. Street, N. W.  
Washington, D. C.

Secretary Ship Structure Committee  
U. S. Coast Guard Headquarters  
1300 E Street, N. W.  
Washington, D. C.

Commander  
Military Sea Transportation Service  
Department of the Navy  
Washington 25, D. C.

U. S. Maritime Administration  
GAO Building  
441 G Street, N. W.  
Washington, D. C.  
Attn: Division of Ship Design  
Division of Research

Superintendent  
U. S. Merchant Marine Academy  
Kings Point, Long Island, New York  
Attn: Capt. L. S. McCready  
(Dept. of Engineering)

Commanding Officer and Director  
U. S. Navy Mine Defense Laboratory  
Panama City, Florida

Commanding Officer  
NROTC and Naval Administrative  
Massachusetts Institute of Technology  
Cambridge 39, Massachusetts

U. S. Army Transportation Research and  
Development Command  
Fort Eustis, Virginia  
Attn: Marine Transport Division

Mr. J. B. Parkinson  
National Aeronautics and Space  
Administration  
1512 H Street, N. W.  
Washington 25, D. C.

Director  
Langley Research Center  
Langley Station  
Hampton, Virginia  
Attn: Mr. I. E. Garrick  
Mr. D. J. Marten

Director Engineering Sciences Division  
National Science Foundation  
1951 Constitution Avenue, N. W.  
Washington 25, D. C.

Director  
National Bureau of Standards  
Washington 25, D. C.  
Attn: Fluid Mechanics Division  
(Dr. G. B. Schubauer)  
Dr. G. H. Keulegan  
Dr. J. M. Franklin

Defense Documentation Center  
Cameron Station  
Alexandria, Virginia (20)  
Office of Technical Services  
Department of Commerce  
Washington 25, D. C.

California Institute of Technology  
Pasadena 4, California  
Attn: Professor M. S. Plesset  
Professor T. Y. Wu  
Professor A. J. Acosta

University of California  
Department of Engineering  
Los Angeles 24, California  
Attn: Dr. A. Powell

Director  
Scripps Institute of Oceanography  
University of California  
La Jolla, California

Professor M. L. Albertson  
Department of Civil Engineering  
Colorado A and M College  
Fort Collins, Colorado

Professor J. E. Cermak  
Department of Civil Engineering  
Colorado State University  
Fort Collins, Colorado

Professor W. R. Sears  
Graduate School of Aeronautical Engineering  
Cornell University  
Ithaca, New York

State University of Iowa  
Iowa Institute of Hydraulic Research  
Iowa City, Iowa  
Attn: Dr. H. Rouse  
Dr. L. Landweber

Massachusetts Institute of Technology  
Cambridge 39, Massachusetts  
Attn: Department of Naval Architecture  
and Marine Engineering  
Professor A. T. Ippen

Harvard University  
Cambridge 38, Massachusetts  
Attn: Professor G. Birkhoff  
(Dept. of Mathematics)  
Professor G. F. Carrier  
(Dept. of Mathematics)

University of Michigan  
Ann Arbor, Michigan  
Attn: Professor R. B. Couch  
(Dept. of Naval Architecture)  
Professor W. W. Willmarth  
(Aero. Engineering Department)

Dr. L. G. Straub, Director  
St. Anthony Falls Hydraulic Laboratory  
University of Minnesota  
Minneapolis 14, Minnesota  
Attn: Mr. J. N. Wetzel  
Professor B. Silberman

Professor J. J. Foody  
Engineering Department  
New York State University Maritime College  
Fort Schuyler, New York

New York University  
Institute of Mathematical Sciences  
25 Waverly Place  
New York 3, New York  
Attn: Professor J. Keller  
Professor J. J. Stoker

The Johns Hopkins University  
Department of Mechanical Engineering  
Baltimore 18, Maryland  
Attn: Professor S. Corrsin  
Professor O. M. Phillips (2)

Massachusetts Institute of Technology  
Department of Naval Architecture and  
Marine Engineering  
Cambridge 39, Massachusetts  
Attn: Professor M. A. Abkowitz, Head

Dr. G. F. Wislicenus  
Ordnance Research Laboratory  
Pennsylvania State University  
University Park, Pennsylvania  
Attn: Dr. M. Sevik

Professor R. C. DiPrima  
Department of Mathematics  
Rensselaer Polytechnic Institute  
Troy, New York

Director  
Woods Hole Oceanographic Institute  
Woods Hole, Massachusetts

Stevens Institute of Technology  
Davidson Laboratory  
Castle Point Station  
Hoboken, New Jersey  
Attn: Mr. D. Savitsky  
Mr. J. P. Breslin  
Mr. C. J. Henry  
Mr. S. Tsakonas

Webb Institute of Naval Architecture  
Crescent Beach Road  
Glen Cove, New York  
Attn: Professor E. V. Lewis  
Technical Library

Executive Director  
Air Force Office of Scientific Research  
Washington 25, D. C.  
Attn: Mechanics Branch

Commander  
Wright Air Development Division  
Aircraft Laboratory  
Wright-Patterson Air Force Base, Ohio  
Attn: Mr. W. Mykytow, Dynamics  
Branch

Cornell Aeronautical Laboratory  
4455 Genesee Street  
Buffalo, New York  
Attn: Mr. W. Targoff  
Mr. R. White

Massachusetts Institute of Technology  
Fluid Dynamics Research Laboratory  
Cambridge 39, Massachusetts  
Attn: Professor H. Ashley  
Professor M. Landahl  
Professor J. Dugundji

Hamburgische Schiffbau-Versuchsanstalt  
Bramfelder Strasse 164  
Hamburg 33, Germany  
Attn: Dr. H. Schwanecke  
Dr. H. W. Lerbs

Institut fur Schiffbau der  
Universitat Hamburg  
Berliner Tor 21  
Hamburg 1, Germany  
Attn: Prof. G. P. Weinblum,

Transportation Technical Research Institute  
1-1057, Mejiro-Cho, Toshima-Ku  
Tokyo, Japan

Max-Planck Institut fur Stromungsforschung  
Bottingerstrasse 6/8  
Gottingen, Germany  
Attn: Dr. H. Reichardt

Hydro-og Aerodynamisk Laboratorium  
Lyngby, Denmark  
Attn: Professor Carl Prohaska

Shipsmodelltanken  
Trondheim, Norway  
Attn: Professor J. K. Lunde  
Versuchsanstalt fur Wasserbau and  
Schiffbau  
Schleuseninsel im Tiergarten  
Berlin, Germany  
Attn: Dr. S. Schuster, Director  
Dr. Grosse

Technische Hogeschool  
Institut voor Toegepaste Wiskunde  
Julianalaan 132  
Delft, Netherlands  
Attn: Professor R. Timman

Bureau D'Analyse et de Recherche  
Appliquees  
47 Avenue Victor Bresson  
Issy-Les-Moulineaux  
Seine, France  
Attn: Professor Siestrunk

Netherlands Ship Model Basin  
Wageningen, The Netherlands  
Attn: Dr. Ir. J. D. vanManen

National Physical Laboratory  
Teddington, Middlesex, England  
Attn: Mr. A. Silverleaf, Superintendent  
Ship Division  
Head, Aerodynamics Division

Head, Aerodynamics Department  
Royal Aircraft Establishment  
Farnborough, Hants, England  
Attn: Mr. M. O. W. Wolfe

Dr. S. F. Hoerner  
148 Busteed Drive  
Midland Park, New Jersey

Boeing Airplane Company  
Seattle Division  
Seattle, Washington  
Attn: Mr. M. J. Turner

Electric Boat Division  
General Dynamics Corporation  
Groton, Connecticut  
Attn: Mr. Robert McCandliss

General Applied Sciences Labs., Inc.  
Merrick and Stewart Avenues  
Westbury, Long Island, New York  
Gibbs and Cox, Inc.  
21 West Street  
New York, New York

Lockheed Aircraft Corporation  
Missiles and Space Division  
Palo Alto, California  
Attn: R. W. Kermeen

Grumman Aircraft Engineering Corp.  
Bethpage, Long Island, New York

Attn: Mr. E. Baird  
Mr. E. Bower  
Mr. W. P. Carl

Midwest Research Institute  
425 Volker Blvd.

Kansas City 10, Missouri

Attn: Mr. Zeydel

Director, Department of Mechanical  
Sciences

Southwest Research Institute  
8500 Culebra Road

San Antonio 6, Texas

Attn: Dr. H. N. Abramson

Mr. G. Ransleben

Editor, Applied Mechanics  
Review

Convair

A Division of General Dynamics  
San Diego, California

Attn: Mr. R. H. Oversmith

Mr. H. T. Brooke

Hughes Tool Company

Aircraft Division

Culver City, California

Attn: Mr. M. S. Harned

Hydronautics, Incorporated

Pindell School Road

Howard County

Laurel, Maryland

Attn: Mr. Phillip Eisenberg

Rand Development Corporation

13600 Deise Avenue

Cleveland 10, Ohio

Attn: Dr. A. S. Iberall

U. S. Rubber Company

Research and Development Department

Wayne, New Jersey

Attn: Mr. L. M. White

Technical Research Group, Inc.

Route 110

Melville, New York, 11749

Attn: Mr. Jack Kotik

Mr. C. Wigley

Flat 102

6-9 Charterhouse Square

London, E. C. 1, England

AVCO Corporation

Lycoming Division

1701 K Street, N. W.

Apt. No. 904

Washington, D. C.

Attn: Mr. T. A. Duncan

Mr. J. G. Baker

Baker Manufacturing Company

Evansville, Wisconsin

Curtiss-Wright Corporation Research  
Division

Turbomachinery Division

Quehanna, Pennsylvania

Attn: Mr. George H. Pedersen

Dr. Blaine R. Parkin

AiResearch Manufacturing Corporation

9851-9951 Sepulveda Boulevard

Los Angeles 45, California

The Boeing Company

Aero-Space Division

Seattle 24, Washington

Attn: Mr. R. E. Bateman

(Internal Mail Station 46-74)

Lockheed Aircraft Corporation

California Division

Hydrodynamics Research

Burbank, California

Attn: Mr. Bill East

National Research Council

Montreal Road

Ottawa 2, Canada

Attn: Mr. E. S. Turner

The Rand Corporation

1700 Main Street

Santa Monica, California

Attn: Technical Library

Stanford University

Department of Civil Engineering

Stanford, California

Attn: Dr. Byrne Perry

Dr. E. Y. Hsu

Dr. Hirsh Cohen

IBM Research Center

P. O. Box 218

Yorktown Heights, New York

Mr. David Wellinger

Hydrofoil Projects

Radio Corporation of America

Burlington, Massachusetts

Food Machinery Corporation

P. O. Box 367

San Jose, California

Attn: Mr. G. Tedrew

Dr. T. R. Goodman

Oceanics, Inc.

Technical Industrial Park

Plainview, Long Island, New York

Professor Brunelle  
Department of Aeronautical Engineering  
Princeton University  
Princeton, New Jersey

Commanding Officer  
Office of Naval Research Branch Office  
230 N. Michigan Avenue,  
Chicago 1, Illinois

University of Colorado  
Aerospace Engineering Sciences  
Boulder, Colorado  
Attn: Prof. M. S. Uberoi

The Pennsylvania State University  
Dept. of Aeronautical Engineering  
Ordnance Research Laboratory  
P. O. Box 30  
State College, Pennsylvania  
Attn: Professor J. William Holl

Institut fur Schiffbau der Universitat Hamburg  
Lammersbeth 90  
2 Hamburg 33, Germany  
Attn: Dr. O. Grim

Technische Hogeschool  
Laboratorium voor Scheepsbouwkunde  
Mekelweg 2, Delft, Netherlands  
Attn: Professor Ir. J. Gerritsma